
| RESEARCH ARTICLE**A Molecular Investigation for *Staphylococcus aureus* in Diabetic Foot Ulcers****Sagia Abbas Gibar Aifari***Department of Basic Sciences, College of Dentistry, University of Wasit, Wasit, Iraq***Corresponding Author:** Sagia Abbas Gibar Aifari, **E-mail:** saifari@uowasit.edu.iq

| ABSTRACT

In Iraq, the high incidence and recurrence rates of diabetic foot ulcers (DFUs) underscore the need for a multifaceted approach to prevention and management. The study aims to direct molecularly detection of the prevalence of *S. aureus* among DFUs patients using molecular assay and then nucleotide sequencing of some local *S. aureus* isolates to be documented in the NCBI and identify its association with the GenBank-BLAST *S. aureus* isolates. Distribution of infection among the age and sex of the study population in addition to the type of medication(s) was also aimed. After disinfecting the skin surrounding the ulcer, pus samples were collected from 73 DFU patients using sterile swabs that were examined molecularly by the PCR assay. DNAs of some positive samples were sequenced and analysed phylogenetically. Targeting the *16S rRNA* gene, 20.55% of DFU swabs had positive reactivity to *S. aureus*. Concerning age, significantly higher prevalence and risk of *S. aureus* were shown in patients aged ≥ 61 years old and to a lesser extent among 51- 60 years old when compared to other age groups: 41-50 and ≤ 40 years old. For sex, although no significant variation was seen among females and males, risk values revealed that males were at higher risk of *S. aureus* infection than females. Regarding the type of medication, DFU patients who received injectable medication have significantly higher values of prevalence rate and risks than those of oral and mixed medications. In relation to phylogeny, the sequenced data of six local *S. aureus* strains were submitted in the NCBI database under specified names (DFU1-DFU6) and GenBank IDs (PQ318365.1- PQ318370.1). The findings of phylogenetic tree analysis, homology sequence identity, and NCBI MSA viewer demonstrated that the local *S. aureus* strains were related to the NCBI-BLAST Iraqi isolate (LC576397.1) at an identity rate ranged from 98.65% to 99.79% and mutation/change at 0.02%. This study confirms, for the first time in Iraq, the phylogenetic linkage of *S. aureus* isolated from DFU patients to the NCBI-GenBank *S. aureus* isolates. This study also shows significant differences in prevalence rates and risks of *S. aureus* among various age and sex groups as well as among DFU patients who received different medications. Therefore, establishing the underlying causes of diabetes and developing effective interventions to prevent and manage its associated complications are crucial steps toward reducing the burden of this widespread disease. Additionally, the high incidence and recurrence rates of DFU underscore the need for a multifaceted approach to prevention and management.

| KEYWORDS

Polymerase chain reaction (PCR), Sequencing analysis, Diabetes mellitus, National Centre For Biotechnology Information (NCBI), Iraq

| ARTICLE INFORMATION**ACCEPTED:** 19 February 2025**PUBLISHED:** 16 March 2025**DOI:** 10.61424/ijmhr.v3.i1.234

1. Introduction

Staphylococcus aureus is an opportunistic Gram-positive, facultative anaerobic coccus that is a common inhabitant of human microbiota, particularly in the nasal passages and on the skin (Ibraheim et al., 2023a). The ubiquitous existence and ability to produce a diverse array of virulence factors (toxins, enzymes, and cell surface proteins) enable it to evade host defenses and cause a wide range of infections that range from relatively mild skin and soft tissue infections to life-threatening conditions such as pneumonia, endocarditis, osteomyelitis, and sepsis (Park and Ronholm, 2021; Ibraheim et al., 2023b). Additionally, the bacterium has the ability to grow in a wide range of

temperatures, pH, and sodium chloride concentrations and to survive dry and stressful environments, which further contributes to causing food-related outbreaks (Alreshidi et al., 2020). In the last decades, the emergence of antibiotic-resistant strains, such as methicillin-resistant *S. aureus*, has further exacerbated the clinical challenge posed by this pathogen, making it a significant public health concern (Abebe and Birhanu, 2023).

Diabetes mellitus is a chronic metabolic disorder characterized by elevated blood glucose levels due to the inability of the body to properly produce (type 1) or utilize (type 2) insulin, a hormone essential for regulating blood sugar (Al-Shaeli et al., 2022). The sustained hyperglycemia associated with diabetes can lead to a multitude of debilitating microvascular and macrovascular complications, affecting the cardiovascular system, kidneys, eyes, and peripheral nerves (Rangel et al., 2019; Mota et al., 2020). Diabetic foot ulcers (DFUs) are a significant complication of diabetes mellitus, causing substantial morbidity, reduced quality of life, and increased healthcare costs (Edmonds et al., 2021). DFUs occur due to a complex interplay between several factors, including peripheral neuropathy, peripheral arterial disease, and impaired wound healing (Davis et al., 2018). Neuropathy leads to loss of sensation and can result in unnoticed trauma and the development of skin lesions (Pokhriyal et al., 2018). Peripheral arterial disease reduces blood flow to the lower extremities, impairing nutrient and oxygen delivery to the wound site, thereby hindering healing (Soyoye et al., 2021). Additionally, diabetes-induced alterations in the inflammatory response and impaired angiogenesis contribute to the delayed and dysfunctional healing characteristic of DFUs (Nirenjen et al., 2023). Hence, the prevention and early detection of these risk factors are crucial in reducing the incidence of DFUs (Lim et al., 2017). In addition, infection control can effectively manage DFUs since several infections lead to sepsis, gangrene, amputation, and even death (Pitocco et al., 2019).

In Iraq, few studies have investigated the prevalence of different microorganisms in DFUs individuals using traditional techniques (Alkhudhairi and Al-Shammari, 2020; Qadir et al., 2020) or by the molecular PCR assay only (Hussein and Saleh, 2024). This study was performed to directly detect the prevalence of *S. aureus* among the DFUs patients using PCR assay and then nucleotide sequencing of some local *S. aureus* isolates to be documented in the NCBI and identify its association with the GenBank-BLAST *S. aureus* isolates. Distribution of infection among the age and sex of the study population in addition to the type of medication(s) was also aimed.

2. Materials and methods

2.1 Ethical approval

The final acceptance and approval were awarded by the Scientific Committee of the Department of Basic Sciences in the College of Dentistry (University of Wasit, Wasit-Iraq). Samples were obtained post-oral agreement of the DFUs patients.

2.2 Samples

After disinfecting the skin surrounding an ulcer, pus samples were collected from 73 DFU's patients using sterile swabs by a circular motion, kept into plastic containers, transported cooled to the laboratory, and subjected directly for molecular processing. Data concerned with the age and sex of study patients and the type of medication(s) were recorded.

2.3 Molecular PCR assaying

According to manufacturers' instructions of the Presto Mini gDNA Bacteria Kit (Geneaid, Taiwan), DNAs were extracted from the pus samples and examined by the Nanodrop System (Thermo Scientific, UK). Targeting the 16S *rRNA* gene, one set of primers [(F: 5'- GGA ACT GAG ACA CGG TCC AG -3') and R: (5'- GGG TCC CCG TCA ATT CCT TT -3')] were designed for the current study based on the NCBI-GenBank *S. aureus* (PQ010750.1) isolate to prepare the Mastermix tubes at a final volume of 20µl using the GoTaq Green Master Mix Kit (Promega, Korea). For PCR reaction, the following conditions of the Thermal Cycler system (BioRad, USA) were followed: 1 cycle for initial denaturation (95°C / 5 minutes); 30 cycles for denaturation (95°C / 30 seconds), annealing (54°C / 30 seconds) and extension (72°C / 30 seconds); and 1 cycle for final extension (72°C / 7 minutes). Electrophoresis of PCR products in Agarose-gel (1.5%) stained with Ethidium Bromide was conducted at 80Am and 100V for 90 minutes, and the product size of positive samples was identified under the UV illuminator (Clinx Science, China) at approximately 616bp.

2.4 Phylogenetic analysis

DNAs of some positive *S. aureus* samples were selected and sent to the Macrogen Company (Korea). The received sequence data were submitted initially in the NCBI-GenBank database to get the specific access numbers for the local *S. aureus* study isolates and then subjected to the multiple sequence alignment analysis and phylogenetic tree analysis in the MEGA-11 Software to identify its identity to the NCBI-BLAST *S. aureus* isolates.

2.5 Statistical analysis

The *t*-test, Odds ratio (OR), as well as Relative Risk (RR) in the GraphPad Prism Software (version 8.0.2) were served to evaluate significant differences between study values at $p < 0.05$, $p < 0.01$, $p < 0.001$, and $p < 0.0001$. The calculated OR involved the value and 95%CI, while the calculation of RR included the value, 95%CI, and NNT (Gharban, 2023).

3. Results

Targeting the 16S rRNA gene, 20.55% (15/73) of study samples showed a positive reactivity to *S. aureus* using the PCR assay (Figures 1, 2).




Figure (1): Molecular prevalence rate (%) of *S. aureus* among the DFU study patients




Figure (2): Agarose-gel electrophoresis of PCR products at 80Am and 100Volt for 90 minutes

Lane (M): Ladder marker (100-1500 bp)

Lanes (2, 3, 4, 8, 10, 11, 14, 15, 18, 19, 20, 23, 27, 28, and 30): Negative samples

Lanes (1, 5, 6, 7, 9, 12, 13, 16, 17, 21, 22, 24, 25, 26, and 29): Positive samples to *S. aureus* at 616bp

Concerning the age of study population, significantly higher values of prevalence rate, OD and RR of *S. aureus* were shown in ≥ 61 years old DFU patients (33.33%, 2.5556, and 1.7778, respectively) and to less extent in 51- 60 years old (23.53%, 1.4066, and 1.2117, respectively) when compared to other age groups; 41-50 years old (8.33%, 0.3052, and 0.4121, respectively) and ≤ 40 years old (0%, 0.2190, and 0.2581, respectively).

For sex, although no significant variation ($p \leq 0.93$) was seen in the prevalence rate of *S. aureus* among females (20%) and males (20.59%), values of OR and RR revealed that DFU males (1.0370 and 1.0244, respectively) were at higher risk of *S. aureus* infection than females (0.9643 and 0.9762, respectively).

Regarding the type of medication, DFU patients who received injectable medication have significantly higher values of prevalence rate (29.03%), OR (2.4545), and RR (1.8) than those of oral (12.5%, 0.4835, and 0.5983, respectively) and mixed (15.39%, 0.5950, and 0.7030, respectively) medications (Table 1).

Table (1): Distribution of *S. aureus* infection among various risk factors

Factor	Group	Total No.	Positive [No. (%)]	OR		RR		
				Value	95%CI	Value	95%CI	NNT
Age (Year)	≤ 40	9	0 (0%)	0.2190	0.0121 to 3.9701	0.2581	0.0167 to 3.9903	6.957 Benefit
	41-50	12	1 (8.33%)	0.3052	0.0362 to 2.5742	0.4121	0.0591 to 2.8718	9.112 Benefit
	51- 60	34	8 (23.53%)	1.4066	0.4504 to 4.3927	1.2517	0.4968 to 3.1539	26.108 Harm
	≥ 61	18	6 (33.33%) *	2.5556 ****	0.7600 to 8.5936	1.7778 ****	0.7082 to 4.4625	9.143 Harm
<i>p-value</i>			<i>0.0118</i>	<i>0.0001</i>	-	<i>0.0001</i>	-	-
Sex	Female	5	1 (20%)	0.9643	0.0997 to 9.3227	0.9762	0.1532 to 6.2191	246 Benefit
	Male	68	14 (20.59%)	1.0370 ***	0.1073 to 10.0261	1.0244 ***	0.1608 to 6.5262	246 Harm
<i>p-value</i>			<i>0.093</i>	<i>0.0002</i>	-	<i>0.0003</i>	-	-
Medication	Oral	16	2 (12.5%)	0.4835	0.0971 to 2.4084	0.5983	0.1482 to 2.4158	13.404 Benefit
	Injection	31	9 (29.03%) *	2.4545 ****	0.7685 to 7.8394	1.8 ****	0.7004 to 4.6262	10 Harm
	Mixed	26	4 (15.39%)	0.5950	0.1686 to 2.1006	0.7030	0.2445 to 2.0213	17.755 Benefit
<i>p-value</i>			<i>0.0458</i>	<i>0.0001</i>	-	<i>0.0001</i>	-	-

In relation to phylogeny, the sequenced data of six local *S. aureus* strains were submitted in the NCBI database under specified names (DFU1-DFU6) and GenBank IDs (PQ318365.1- PQ318370.1). The findings of phylogenetic tree analysis, homology sequence identity, and NCBI MSA viewer demonstrated that the local *S. aureus* strains were related to the NCBI-BLAST Iraqi isolate (LC576397.1) at an identity rate ranged from 98.65% to 99.79% and mutation/change at 0.02% (Figures 3-5, Table 2).




Figure (4): Phylogenetic tree analysis of the local and NCBI-BLAST *S. aureus* isolates/strains using MEGA-11 software




Figure (5): Multiple sequence alignment of the local and NCBI-BLAST *S. aureus* isolates/strains using NCBI MSA Viewer

Table (2): Homology Sequence identity (%) for local and NCBI-BLAST *S.aureus* isolates

Local isolate		NCBI isolate			
Name	Access No.	Species	Country	Access No.	%
DFU1	PQ318365.1	<i>S. aureus</i>	Iraq	LC576397.1	99.72
DFU2	PQ318366.1	<i>S. aureus</i>	Iraq	LC576397.1	99.51
DFU3	PQ318367.1	<i>S. aureus</i>	Iraq	LC576397.1	99.79
DFU4	PQ318368.1	<i>S. aureus</i>	Iraq	LC576397.1	98.66
DFU5	PQ318369.1	<i>S. aureus</i>	Iraq	LC576397.1	98.65
DFU6	PQ318370.1	<i>S. aureus</i>	Iraq	LC576397.1	99.68

4. Discussion

Worldwide, several risk factors have been identified to predispose individuals with diabetes to the development of foot ulcers. These include poor glycemic control, longer duration of diabetes, older age, obesity, smoking, and a history of previous foot ulceration or amputation. Foot deformities, such as Charcot's foot, and inadequate foot care can also increase the risk of ulceration (Ahmad, 2016; Ntuli et al., 2018; Lin et al., 2020).

Different studies demonstrated that *S. aureus* was the more predominant Gram-positive pathogen responsible for infections of mild-to-moderate severity in diabetic patients with foot ulcers (Esposito et al., 2008; Bengalorkar, 2011; Appapalam et al., 2021). Dunyach-Remy et al. (2016) described the role of *S. aureus* in DFUs and the implication of its toxins in the establishment of infection. In this study, PCR assay demonstrated the presence of *S. aureus* infection in ulcers of 20.55% DFU patients, which in agreement with that reported previously in India (24.5%) by Shankar et al. (2005) and recently in Iraq by Hussein and Saleh (2024) who identified *S. aureus* in 24.28% of the DFU patients attended to two hospitals in Alsamawa and Baghdad provinces during the 2022s. In comparison to other global studies, a study conducted by Tentolouris et al. (2006) in Greece has detected 18 out of 91-foot ulcers were infected by *S. aureus* in 19.78%, which, methicillin-resistant *S. aureus* (MRSA) isolates were more common among patients with infected foot ulcers (61.1%). Eleftheriadou et al. (2010) showed that the prevalence rate of MRSA in DFUs was 15-30%, and there is an alarming trend for increase in many countries. In 2014, a study conducted in the USA characterized 25 different *S. aureus* isolates in DFU patients, and most of these isolates resembled genetically those isolated from patients with atopic dermatitis (Vu et al., 2014). Shettigar et al. (2016) detected that 43% (86/200) of DFUs were infected with *S. aureus*, and virulence genes were found in 94.19% (81/86) isolates, with the presence of more than one gene in 41 isolates. In a cross-sectional study among 83 adult DFU patients in Kenya, the findings showed that 94% of swab samples were positive to Gram-negative (65%) and Gram-positive (29%) with detection of *S. aureus* in 16% of these isolates. In Iraq, the culture findings of Qadir et al. (2020) observed that the prevalence rate of Gram-positive bacteria was significantly greater than that of Gram-negative bacteria and that *S. aureus* encompasses 54.7% of identified Gram-positive isolates. Another study recorded that 201 out of 863 (23.29%) DFU swabs were infected with *S. aureus*, in which MRSA isolates were detected in 15.4% (Stańkowska et al., 2022). In a recent systemic review and meta-analysis encompassed 40 studies across 20 countries, Zhou et al. (2024) found that the overall prevalence of MRSA in DFU was 17% with significant regional variation: 61% in South America, 20% in North America, 19% in Europe, 13% in Africa and 11% in others. However, the uncontrolled use of antibiotics in both hospitalized and non-hospitalized patients may contribute effectively to increasing the prevalence and risk of *S. aureus* among DFU patients.

The current study shows that the variable prevalence of *S. aureus* among different ages, sexes, and medications is high. The fact that more *S. aureus* isolates were seen in DFU patients older than 60 years may be due to either the high incidence of chronic diseases or attenuation of immunity in advanced age. Also, various factors such as professional activities and lifestyle may cause the feet to tolerate more pressure, increasing the incidence of

consequences that are more common in older people. The main results of Cervantes-García et al. (2015) included 34% positive *S. aureus* isolates and indicated that the grade-4 of DFU is more common in patients aged 54-65 and 66-88 years old and having diabetes for 11-15 years or longest period (16-30 years). Subsequently, the authors found no significant variation in the prevalence of different microbes isolated from DFU lesions but showed that the majority had localized gangrene, and the majority of women had deep ulcers with cellulitis and/or abscesses in the skin. Mohammed et al. (2016) mentioned that older age, long diabetes duration, multiple anti-diabetics, hard physical activity, and bad foot care are the common risk factors for the development of DFUs. Lee et al. (2017) showed that the MRSA infection did not differ between DFU patients of both age groups, ≥ 65 (14.7%) and < 65 (15.8%) years, as well as between males (15%) and females (16.1%). In an Iraqi study, Qadir et al. (2020) identified that DFU was more common in individuals aged 50-59 years and, to a lesser extent, in 60-69 and ≥ 70 years old when compared to < 40 and 40-49 years old. Also, they recorded that DFU is more related to oral drugs than insulin and mixed (oral drugs and insulin) medications. A cross-sectional study conducted by Anafo et al. (2021) found that the prevalence rate of *S. aureus* among DFU patients was 19%, and infection is more common in advanced ages, 30-60 years (42%) and > 60 years (57%), as well as in females (54%) more than males (46%). Among a total of 1771 patients, Vanherwegen et al. (2023) reported that the vast majority of DFU infection was seen in males (72%), and lesions in males being deeper, more frequently, displaying probe to bone and more frequently deeply infected, while women had a 265 higher chance of healing without major amputation as a first event.

Staphylococcus aureus is characterized by both vertical and horizontal evolution, and the vertical historical signal is often strong enough to make phylogenetic reconstruction meaningful and useful (Planet et al., 2017). In the current study, phylogenetic analysis revealed that the local *S. aureus* strains were highly identical to the NCBI-BLAST *S. aureus* strains isolated from eye infection of humans at Basra province (Iraq) and submitted in the NCBI-GenBank in 2020. These findings support the fact that *S. aureus* can cause different skin and skin structure infections, including DFUs, osteoarticular and ocular infections, pneumonia, and endocarditis (Isah, 2021; Van Dorpe, 2021). However, genotyping methods add precision to the identification of *S. aureus*, and partial 16S rRNA sequencing can be served largely in the identification of different pathogenic isolates/strains (Kosecka-Strojek et al., 2019; Wilson et al., 2019).

5. Conclusion

This study confirms, for the first time in Iraq, the phylogenetic linkage of *S. aureus* isolated from DFU patients to the NCBI-GenBank *S. aureus* isolates. This study also shows significant differences in prevalence rates and risks of *S. aureus* among various age and sex groups as well as among DFU patients who received different medications. Therefore, establishing the underlying causes of diabetes and developing effective interventions to prevent and manage its associated complications are crucial steps toward reducing the burden of this widespread disease. Additionally, the high incidence and recurrence rates of DFU underscore the need for a multifaceted approach to prevention and management.

Acknowledgment: The author thanks all workers and patients who contributed to completing this work through sample collection.

Conflicts of interests: The author declares no conflict of interest.

Funding: This study was privately funded.

References

- [1] Abebe, A.A., and Birhanu, A.G. (2023). Methicillin resistant *Staphylococcus aureus*: molecular mechanisms underlying drug resistance development and novel strategies to Combat. *Infection and Drug Resistance*, 7641-7662.
- [2] Ahmad, J. (2016). The diabetic foot. *Diabetes and Metabolic Syndrome: Clinical Research and Reviews*, 10(1), 48-60.
- [3] Alkhdhairy, M.K., and Al-Shammari, M.M.M. (2020). Prevalence of metallo- β -lactamase-producing *Pseudomonas aeruginosa* isolated from diabetic foot infections in Iraq. *New microbes and new infections*, 35, 100661.
- [4] Alreshidi, M.M., Dunstan, R.H., Macdonald, M.M., Gottfries, J., and Roberts, T.K. (2020). The uptake and release of amino acids by *Staphylococcus aureus* at mid-exponential and stationary phases and their corresponding responses to changes in temperature, pH and osmolality. *Frontiers in Microbiology*, 10, 3059.

- [5] Al-Shaeli, S.J., Hussen, T.J., and Ethaeb, A.M. (2022). Effect of honey bee venom on the histological changes of testes and hormonal disturbance in diabetic mice. *Veterinary World*, 15(9), 2357.
- [6] Anafo, R.B., Atiase, Y., Dayie, N.T., Kotey, F.C., Tetteh-Quarcoo, P.B., Duodu, S., and Donkor, E.S. (2021). Methicillin-resistant *Staphylococcus aureus* (MRSA) infection of diabetic foot ulcers at a tertiary care hospital in Accra, Ghana. *Pathogens*, 10(8), 937.
- [7] Appapalam, T.S., Muniyan, A., Vasanthi Mohan, K., and Panchamoorthy, R. (2021). A study on isolation, characterization, and exploration of multiantibiotic-resistant bacteria in the wound site of diabetic foot ulcer patients. *The International Journal of Lower Extremity Wounds*, 20(1), 6-14.
- [8] Bengalorkar, G.M. (2011). Diabetic foot infections—A review. *Int J Biol Med Res*, 2(1), 453-460.
- [9] Cervantes-García, E., García-González, R., Reséndiz-Albor, A., and Salazar-Schettino, P.M. (2015). Infections of diabetic foot ulcers with methicillin-resistant *Staphylococcus aureus*. *The International Journal of Lower Extremity Wounds*, 14(1), 44-49.
- [10] Davis, F.M., Kimball, A., Boniakowski, A., and Gallagher, K. (2018). Dysfunctional wound healing in diabetic foot ulcers: new crossroads. *Current diabetes reports*, 18, 1-8.
- [11] Dunyach-Remy, C., Ngba Essebe, C., Sotto, A., and Lavigne, J.P. (2016). *Staphylococcus aureus* toxins and diabetic foot ulcers: role in pathogenesis and interest in diagnosis. *Toxins*, 8(7), 209.
- [12] Edmonds, M., Manu, C., and Vas, P. (2021). The current burden of diabetic foot disease. *Journal of clinical orthopaedics and trauma*, 17, 88-93.
- [13] Eleftheriadou, I., Tentolouris, N., Argiana, V., Jude, E., and Boulton, A.J. (2010). Methicillin-resistant *Staphylococcus aureus* in diabetic foot infections. *Drugs*, 70, 1785-1797.
- [14] Esposito, S., Leone, S., Noviello, S., Fiore, M., Ianniello, F., Felaco, F.M., and Sarli, E. (2008). Foot infections in diabetes (DFIs) in the out-patient setting: an Italian multicentre observational survey. *Diabetic medicine*, 25(8), 979-984.
- [15] Gharban, H.A. (2023). Molecular prevalence and phylogenetic confirmation of bovine trichomoniasis in aborted cows in Iraq. *Veterinary world*, 16(3), 580-587.
- [16] Hussein, S.Z., and Saleh, G.M. (2024). Molecular Detection of Virulence Factors Genes for *Staphylococcus aureus* in Diabetic Foot Ulcers in Iraq. *Ibn AL-Haitham Journal For Pure and Applied Sciences*, 37(3), 98-105.
- [17] Ibraheim, H.K., Madhi, K.S., Baqer, G.K., and Gharban, H.A. (2023a). Effectiveness of raw bacteriocin produced from lactic acid bacteria on biofilm of methicillin-resistant *Staphylococcus aureus*. *Veterinary World*, 16(3), 491-499.
- [18] Ibraheim, H.K., Fayez, R.A., Jasim, A.S., and Gharban, H.A. (2023b). Role of nuc gene in *Staphylococcus aureus* to phagocytic activity in different cattle infections. *Open Veterinary Journal*, 13(8), 1021-1026.
- [19] Isah, R.M. (2021). Antibacteria activities of crude *Curcuma longa* extract mediated silver nanoparticles against isolates from diabetic patients with foot infections (Doctoral dissertation).
- [20] Kosecka-Strojek, M., Sabat, A.J., Akkerboom, V., Becker, K., Van Zanten, E., Wisselink, G., and Friedrich, A.W. (2019). Development and validation of a reference data set for assigning *Staphylococcus* species based on next-generation sequencing of the 16S-23S rRNA region. *Frontiers in cellular and infection microbiology*, 9, 278.
- [21] Lee, J.S., Son, S.T., and Han, S.K. (2017). Risk factors of methicillin-resistant *Staphylococcus aureus* and *Pseudomonas* infection in diabetic foot ulcers in Korea. *Journal of Wound Management and Research*, 13(2), 29-34.
- [22] Lim, J.Z.M., Ng, N.S.L., and Thomas, C. (2017). Prevention and treatment of diabetic foot ulcers. *Journal of the Royal Society of Medicine*, 110(3), 104-109.
- [23] Lin, C., Liu, J., and Sun, H. (2020). Risk factors for lower extremity amputation in patients with diabetic foot ulcers: A meta-analysis. *PloS one*, 15(9), e0239236.
- [24] Mohammed, S.I., Mikhael, E.M., Ahmed, F.T., Al-Tukmagi, H.F., and Jasim, A.L. (2016). Risk factors for occurrence and recurrence of diabetic foot ulcers among Iraqi diabetic patients. *Diabetic foot and ankle*, 7(1), 29605.
- [25] Mota, R.I., Morgan, S.E., and Bahnson, E.M. (2020). Diabetic vasculopathy: macro and microvascular injury. *Current pathobiology reports*, 8, 1-14.
- [26] Nirenjen, S., Narayanan, J., Tamilanban, T., Subramaniyan, V., Chitra, V., Fuloria, N.K., and Selvaraj, S. (2023). Exploring the contribution of pro-inflammatory cytokines to impaired wound healing in diabetes. *Frontiers in immunology*, 14, 1216321.
- [27] Ntuli, S., Swart, A., and Lambert, C.V. (2018). Risk factors for diabetic foot ulceration. *South African Journal of Diabetes and Vascular Disease*, 15(1), 32-36.
- [28] Park, S., and Ronholm, J. (2021). *Staphylococcus aureus* in agriculture: lessons in evolution from a multispecies pathogen. *Clinical microbiology reviews*, 34(2), 10-1128.
- [29] Pitocco, D.S.T.S.M., Spanu, T., Di Leo, M., Vitiello, R., Rizzi, A., Tartaglione, L., and Sanguinetti, M. (2019). Diabetic foot infections: a comprehensive overview. *European Review for Medical and Pharmacological Sciences*, 23.
- [30] Planet, P.J., Narechania, A., Chen, L., Mathema, B., Boundy, S., Archer, G., and Kreiswirth, B. (2017). Architecture of a species: phylogenomics of *Staphylococcus aureus*. *Trends in microbiology*, 25(2), 153-166.
- [31] Pokhriyal, V., Kothiyal, P., Kumar, N., and Kaushik, S. (2018). A review on diabetic neuropathy: Complications and treatment. *Asian J Pharm Pharmacol*, 4(4), 413-420.

- [32] Qadir, A.N., Mahmoud, B.M., Mahwi, T.O., Al-Attar, D.M.R.A., and Mahmood, S.O. (2020). Prevalence of microorganisms and antibiotic sensitivity among patients with diabetic foot ulcer in Sulaimani City, Iraq. *Hospital Practices and Research*, 5(2), 56-63.
- [33] Rangel, É.B., Rodrigues, C.O., and De Sa, J.R. (2019). Micro-and macrovascular complications in diabetes mellitus: preclinical and clinical studies. *Journal of Diabetes Research*, 2019.
- [34] Shankar, E.M., Mohan, V., Premalatha, G., Srinivasan, R.S., and Usha, A.R. (2005). Bacterial etiology of diabetic foot infections in South India. *European journal of internal medicine*, 16(8), 567-570.
- [35] Shettigar, K., Jain, S., Bhat, D.V., Acharya, R., Ramachandra, L., Satyamoorthy, K., and Murali, T.S. (2016). Virulence determinants in clinical *Staphylococcus aureus* from monomicrobial and polymicrobial infections of diabetic foot ulcers. *Journal of medical microbiology*, 65(12), 1392-1404.
- [36] Soyoye, D.O., Abiodun, O.O., Ikem, R.T., Kolawole, B.A., and Akintomide, A.O. (2021). Diabetes and peripheral artery disease: A review. *World journal of diabetes*, 12(6), 827.
- [37] Stańkowska, M., Garbacz, K., Korzon-Burakowska, A., Bronk, M., Skotarczak, M., and Szymańska-Dubowik, A. (2022). Microbiological, clinical and radiological aspects of diabetic foot ulcers infected with methicillin-resistant and-sensitive *Staphylococcus aureus*. *Pathogens*, 11(6), 701.
- [38] Tentolouris, N., Petrikos, G., Vallianou, N., Zachos, C., Daikos, G.L., Tsapogas, P., and Katsilambros, N. (2006). Prevalence of methicillin-resistant *Staphylococcus aureus* in infected and uninfected diabetic foot ulcers. *Clinical microbiology and infection*, 12(2), 186-189.
- [39] Van Dorpe, I. (2021). Abstracts presented at the 25th Annual Congress of the Belgian Society of Internal Medicine, 10-11 December 2021, Dolce La Hulpe, La Hulpe, Belgium. *Acta Clinica Belgica*, 76(sup2), 1-68.
- [40] Vanherwegen, A.S., Lauwers, P., Lavens, A., Doggen, K., Dirinck, E., and Initiative for Quality Improvement and Epidemiology in multidisciplinary Diabetic Foot Clinics (IQED-Foot) Study Group. (2023). Sex differences in diabetic foot ulcer severity and outcome in Belgium. *PLoS One*, 18(2), e0281886.
- [41] Vu, B.G., Stach, C.S., Salgado-Pabón, W., Diekema, D.J., Gardner, S.E., and Schlievert, P.M. (2014). Superantigens of *Staphylococcus aureus* from patients with diabetic foot ulcers. *The Journal of infectious diseases*, 210(12), 1920-1927.
- [42] Wilson, D.J., Middleton, J.R., Adkins, P.R., and Goodell, G.M. (2019). Test agreement among biochemical methods, matrix-assisted laser desorption ionization–time of flight mass spectrometry, and 16S rRNA sequencing for identification of microorganisms isolated from bovine milk. *Journal of clinical microbiology*, 57(3), 10-1128.
- [43] Zhou, S., Hu, X., Wang, Y., Fei, W., Sheng, Y., and Que, H. (2024). The Global Prevalence of Methicillin-Resistant *Staphylococcus Aureus* in Patients with Diabetic Foot Ulcers: A Systematic Review and Meta-Analysis. *Diabetes, Metabolic Syndrome and Obesity*, 563-574.